

On auto and hetero Backlund transformation for the Hamilton-Jacobi equations

Alexei Sozonov

Saint Petersburg State University
Universitetskaya nab., 7-9
Saint Petersburg, Russia

On auto and hetero Backlund transformation for the Hamilton-Jacobi equations The primary aim of the study was to investigate Backlund transformations (BT) and to discuss some correspondence between the integrable cases of the Henon-Heiles system. This correspondence may be considered a finite-dimensional counterpart of the hetero Backlund transformation that can be used for building and classifying of integrable systems. An application of the possible finite-dimensional analogues of the well-studied Backlund transformations for nonlinear differential equations to three Henon-Heiles systems is discussed. Using the known 2×2 Lax matrix and special similarity transformation we obtain new unique Lax matrix with the following properties: first off-diagonal element of Lax matrix defines initial parabolic coordinates on the plane $v_{1,2}$; second off-diagonal element has only two commuting and functionally independent zeroes $u_{1,2}$; the conjugated momenta for u and v variables are the values of the diagonal element; Thus we have two families of variables of separation for the first Henon-Heiles system and can explicitly define canonical transformation between them. After that we converted this special auto-BT to some analogue of the hetero-BT which relates different Hamilton-Jacobi equations associated with the three different integrable cases of the Henon-Heiles system.